

Wigner 111, Budapest, Nov 2013

## **Fractional and Majorana Fermions**

R. Jackiw (MIT)

A bit of physics history about Dirac and his equation:

Dirac was looking for a relativistic equation for electrons and eventually arrived at the following first-order matrix equation



Dirac Matrix Equation

$$(\boldsymbol{\alpha} \cdot \mathbf{p} + \beta m)\Psi = i\frac{\partial}{\partial t} \Psi$$

$\Psi = 4$ -component complex function (charged excitations)

$\boldsymbol{\alpha}, \beta \Rightarrow$  four  $4 \times 4$  "Dirac" numerical matrices

$$\mathbf{p} = \frac{1}{i}\nabla, \quad m = \text{mass parameter}$$

Dirac matrices ensure that iteration implies

$$-\frac{\partial^2}{\partial t^2} \Psi = -\nabla^2 \Psi + m^2 \Psi$$

Dirac equation =  $\pm\sqrt{\text{massive wave equation}}$

To expose properties of the Dirac equation, we make the usual decomposition

$$\Psi = e^{-iEt} \psi \Rightarrow [\boldsymbol{\alpha} \cdot \mathbf{p} + \beta m] \psi = E \psi$$

and discover that there exist positive energy solutions  $E > 0$ , which can describe electrons.

But because “square roots” come in both signs, there exist also negative energy solutions,  $E < 0$ . These need interpretation because they cannot describe electrons, which carry positive energy. After some hesitation, Dirac concluded that the negative energy solutions correspond to anti-electrons, *viz.* positrons, which soon were discovered.

In a further conceptual leap, Dirac posited that in the ground state all negative energy levels are filled, but nevertheless the charge of the ground state is zero. This means that the Dirac equation is really a many particle equation, where the particles populate the energy levels.

The Dirac equation is a beautiful equation, with a rich hidden physical structure. It goes beyond a single particle interpretation and it predicts new (anti-)particles: the positrons. These characteristics make it a beautiful equation for physics.

Also it is a beautiful equation for mathematics because by using matrices, it succeeds in taking a square root of a second order differential equation.

The mathematical and physical beauty of the Dirac equation suggests to mathematicians and physicists that deformations of the equation may also yield beautiful and interesting results.

## Which deformations should we consider?

One alteration that can be made is to reconsider the equation in dimensions different from the three spatial and one time dimension. If we take fewer dimensions we gain the mathematical advantage of simplicity and also have the possibility of describing physical systems that are confined to lower dimensions, for example to a line or to a plane. Such configurations can occur in condensed matter physics. There one encounters situations where the low energy dynamics is well described by a matrix equation, linear in the momenta. Also there may be a constant mass term which separates the positive energy solutions from the negative energy ones by a “gap.” Depending on the nature of the material, the equation may describe excitations on a line, on the plane in addition to those in the three-dimensional bulk.

## Dirac Equations (First-order matrix equations)

$$[\boldsymbol{\alpha} \cdot \mathbf{p} + \beta m]\psi = E\psi$$

continuum solutions  $E > |m|$  and  $E < -|m|$

“vacuum”:

Particle interpretation:  $E < 0$  states filled (antiparticles)  
 $E > 0$  states empty (particles)

Condensed matter:  $E < 0$  states filled (valence band)  
interpretation:  $E > 0$  states empty (conduction band)  
 $m$  produces gap  $2|m|$

“vacuum” carries no net charge

One dimension, physics on a line, Dirac matrices realized with  
 $2 \times 2$  Pauli matrices

$$1-d: \quad \alpha = \sigma_2, \quad \beta = \sigma_1, \quad p = \frac{1}{i} \frac{d}{dx}$$

A further more profound deformation allows the mass term  $m$  to depend on position.

### **What sort of dependence should we consider?**


Surely a weak dependence will produce only insignificant changes from the usual homogenous mass case; we are interested in a significant dependence on position, which could significantly alter the physical situation. Note that the gap  $2|m|$  depends only on the magnitude of  $m$ , and not on its sign.  $+m$  produces the same gap as  $-m$ . This suggests a deformation of the mass term that interpolates between positive and negative values. In this way we are led to a Dirac equation in the presence of a defect.

(mass term position-dependent, soliton)


$$m \rightarrow \varphi(\mathbf{r})$$

$$[\boldsymbol{\alpha} \cdot \mathbf{p} + \beta \varphi(\mathbf{r})]\psi = E\psi$$


1-d kink



2-d vortex



3-d magnetic monopole



Find:

continuum solutions  $E > 0, E < 0$

AND isolated, normalizable  $E = 0$  solution

“mid-gap” state is found by explicit calculation

is guaranteed by index theorems

CENTRAL QUESTION: in “vacuum” is mid-gap state empty or filled, what is its charge?

UNEXPECTED ANSWER: charge  $Q = \pm \frac{1}{2}$ .



## Fractional Charge (Analytic Derivation)

Vacuum charge density:

$$\rho(\mathbf{r}) = \int_{-\infty}^0 dE \rho_E(\mathbf{r}) \quad \rho_E = \psi_E^\dagger \psi_E$$

renormalized charge in soliton background

$$Q = \int d\mathbf{r} \int_{-\infty}^{0-} dE (\rho_E^s(\mathbf{r}) - \rho_E^0(\mathbf{r}))$$

Evaluation simple in the presence of an energy reflection symmetry:

$$\rho_{-E} = \rho_E \text{ (charge conjugation)}$$

## Fractional Charge Calculation

Completeness: 
$$\int_{-\infty}^{\infty} dE \psi_E^\dagger(\mathbf{r}) \psi_E(\mathbf{r}') = \delta(\mathbf{r} - \mathbf{r}')$$

$$\Rightarrow \int_{-\infty}^{\infty} dE [\rho_E^s(\mathbf{r}) - \rho_E^0(\mathbf{r})] = 0$$

Conjugation ( $\rho_E = \rho_{-E}$ ) and zero mode  $\Rightarrow$

$$\int_{-\infty}^{0-} dE (2\rho_E^s(\mathbf{r}) - 2\rho_E^0(\mathbf{r})) + \psi_{E=0}^\dagger(\mathbf{r}) \psi_{E=0}(\mathbf{r}) = 0$$

$$\int_{-\infty}^{0-} dE (\rho_E^s(\mathbf{r}) - \rho_E^0(\mathbf{r})) = -\frac{1}{2} \psi_{E=0}^\dagger(\mathbf{r}) \psi_{E=0}(\mathbf{r})$$

$$Q = -\frac{1}{2}$$

Any dimension!

Empty mid-gap state:  $Q = -\frac{1}{2}$

Filled mid-gap state:  $Q = +\frac{1}{2}$

Eigenvalue, not expectation value!

## One Dimensional Example (Polyacetylene)

Dirac equation with varying mass  $[\sigma_2 p + \sigma_1 \varphi(x)]\psi = E\psi$

$$E \leq 0 \quad \left[ \begin{pmatrix} 0 & -\frac{d}{dx} \\ \frac{d}{dx} & 0 \end{pmatrix} + \begin{pmatrix} 0 & \varphi(x) \\ \varphi(x) & 0 \end{pmatrix} \right] \begin{bmatrix} \psi_E^u \\ \psi_E^l \end{bmatrix} = E \begin{bmatrix} \psi_E^u \\ \psi_E^l \end{bmatrix}$$

$$E = 0 \quad \begin{pmatrix} 0 & -\frac{d}{dx} + \varphi(x) \\ \frac{d}{dx} + \varphi(x) & 0 \end{pmatrix} \begin{bmatrix} \psi_0^u \\ \psi_0^l \end{bmatrix} = 0$$

$$\psi_0^{u,l} = N \exp \mp \left[ \int^x dx' \varphi(x') \right]$$

Rebbi & RJ, *PRD* **13**, 3398 (76)

Su, Schrieffer & Heeger, *PRL* **42**, 1698 (79)

[high conductivity in polymers]

## Derivation within Condensed Matter


Polyacetylene Realization of  $(1 - d)$  Dirac Equation

$$H = \sigma_2 p + \sigma_1 m$$


Kinetic term: linearization at Fermi level

$\pm|m|$ : Gap preserving mass term

(due to energetic instability ~ Peierls)




$m \rightarrow \varphi(x)$ : interpolating between  $\pm|m|$



# Graphene Realization of (2-d) Dirac Equation

Graphene hexagonal lattice



Kinetic term:

Linearization at Fermi level

2 “Dirac points” per sublattice

$\Rightarrow 4 \times 4$  Dirac Hamiltonian in 2 spatial dimensions

Wallace, *PR* **71**, 662 (47); Semenoff, *PRL* **53**, 2449 (84);

Gaim & Novoselov, Nobel Prize (10)

Gap inducing term ? [proposal: Jadecola et al. *PRL* **110**,  
176603 (13)]

## Dirac Hamiltonian

$$H = \underbrace{\psi^\dagger \boldsymbol{\alpha} \cdot \mathbf{p} \psi}_{\text{linearization at Fermi level}} + \underbrace{\psi^\dagger \beta [\varphi_{re} - i \varphi_{im} \gamma_5] \psi}_{\text{Kekulé distortion}} \quad (4 \times 4)$$

$$\varphi \equiv \varphi_{re} + i \varphi_{im}$$

$$\boldsymbol{\alpha} = \begin{pmatrix} \boldsymbol{\sigma} & 0 \\ 0 & -\boldsymbol{\sigma} \end{pmatrix}, \quad \beta = \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix}, \quad \gamma_5 = \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix}, \quad \mathbf{p} = \frac{1}{i} \boldsymbol{\nabla}$$

(All vectors are 2-dimensional)

linearization  
at Fermi level

Kekulé distortion  
 $\varphi$  constant:  $\varphi_0 \Rightarrow$  mass gap  
 $\varphi$  soliton/vortex:  $\varphi_s = |\varphi(r)|e^{i\theta}, \varphi(0) = 0, |\varphi(\infty)| = \varphi_0$   
 $\Rightarrow$  mid-gap state

Conclusion :  $Q = \pm \frac{1}{2}$

Hou, Chamon & Mudry, PRL **98**, 186809 (07) [cond-mat/0609740]

## Majorana Equation

electrically charged particles:

particle is different from anti-particle  
described by complex field

electrically neutral particles:

particle can be identified with its anti-particle  
described by real field

e.g. neutral pion ( $S = 0$ )  
photon ( $S = 1$ )  
graviton ( $S = 2$ ) } all bosons

Majorana fermion = neutral fermion (identified with anti-particle)



Majorana Matrix Equation

$$(\boldsymbol{\alpha} \cdot \mathbf{p} + \beta m)\Psi = i \frac{\partial}{\partial t} \Psi$$

$\Psi$  real (neutral excitations)

$$\mathbf{p} = \frac{1}{i} \nabla \text{ imaginary}$$

$$\boldsymbol{\alpha} \text{ real} = \boldsymbol{\alpha}^*$$

$$\beta \text{ imaginary} = -\beta^*$$

} Majorana Representation

## Majorana Representation

$$\alpha_M^1 = \begin{pmatrix} 0 & \sigma_1 \\ \sigma_1 & 0 \end{pmatrix} \quad \alpha_M^2 = \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix} \quad \alpha_M^3 = \begin{pmatrix} 0 & \sigma_3 \\ \sigma_3 & 0 \end{pmatrix}$$

$$\beta_M = \begin{pmatrix} 0 & \sigma_2 \\ \sigma_2 & 0 \end{pmatrix} \quad \Psi_M^* = \Psi_M$$

Majorana in arbitrary representation

$$C\alpha^*C^{-1} = \alpha, \quad C\beta^*C^{-1} = -\beta \quad C\Psi^* = \Psi$$

e.g. Weyl

$$\alpha = \begin{pmatrix} \boldsymbol{\sigma} & 0 \\ 0 & -\boldsymbol{\sigma} \end{pmatrix} \quad \beta = \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix} \quad C = \begin{pmatrix} 0 & -i\sigma_2 \\ i\sigma_2 & 0 \end{pmatrix}$$

$$\begin{pmatrix} \boldsymbol{\sigma} \cdot \mathbf{p} & m \\ m & -\boldsymbol{\sigma} \cdot \mathbf{p} \end{pmatrix} \begin{pmatrix} \psi \\ \chi \end{pmatrix} = i\frac{\partial}{\partial t} \begin{pmatrix} \psi \\ \chi \end{pmatrix}$$

$$C\Psi^* = \Psi \quad \Rightarrow \quad \chi = i\sigma_2 \psi^*$$

$$\boldsymbol{\sigma} \cdot \mathbf{p} \psi + i\sigma_2 m \psi^* = i\frac{\partial}{\partial t} \psi \quad (2 \times 2)$$

$\psi$  mixes with  $\psi^*$

NB  $C = I$  in Majorana representation



## Majorana Equation (2 component)

$$\boldsymbol{\sigma} \cdot \mathbf{p} \psi + i\sigma_2 m \psi^* = i \frac{\partial}{\partial t} \psi$$

NB. Dirac mass term: preserves quantum numbers (charge, particle number)

Majorana mass term: does not preserve any quantum numbers

⇒ no distinction between particle and anti-particle  
since there are no conserved quantities to tell  
them apart, particle is its own anti-particle

Dirac field operator

$$\Psi = \sum_{E>0} \left( a_E e^{-iEt} \Psi_E + b_E^\dagger e^{iEt} C \Psi_E^* \right)$$

Majorana field operator

$$\Psi = \sum_{E>0} \left( a_E e^{-iEt} \Psi_E + a_E^\dagger e^{iEt} C \Psi_E^* \right)$$

# Are there Majorana fermions in Nature?

neutrinos?

– recent development in neutrino physics

experimental observation of neutrino oscillations  $\Rightarrow$


- neutrinos have mass ( $< 0.1\text{eV}$ )
- lepton number is not conserved separately for each flavor.

$\Rightarrow$  they could be Majorana fermions

Hypothetical Majorana fermions:

- supersymmetry – supersymmetric partners of photon, neutral Higgs boson, etc. are necessarily Majorana fermions
- cosmology – dark matter candidates

# Majorana fermions in superconductor in contact with a topological insulator



Hamiltonian density for the model:

$$H = \psi^* \left( \boldsymbol{\sigma} \cdot \frac{1}{i} \nabla - \mu \right) \psi + \frac{1}{2} (\Delta \psi^* i \sigma_2 \psi^* + h.c.)$$

$$\psi = \begin{pmatrix} \psi_{\uparrow} \\ \psi_{\downarrow} \end{pmatrix}, \boldsymbol{\sigma} = (\sigma_1, \sigma_2),$$

$\mu$  is chemical potential,  $\Delta$  is the order parameter

$\Delta$  may be constant:  $\Delta = \Delta_0$

or take vortex profile:  $\Delta(\mathbf{r}) = v(r)e^{i\theta}, v(0) = 0, v(\infty) = \Delta_0.$

Equation of motion:  $i \partial_t \psi = (\boldsymbol{\sigma} \cdot \mathbf{p} - \mu) \psi + \Delta i \sigma_2 \psi^*$

In the absence of  $\mu$ , and with constant  $\Delta$ , the above system is a (2+1)-dimensional version of the (3+1)-dimensional, two component Majorana equation!

$\Rightarrow$  governs chargeless spin  $\frac{1}{2}$  fermions with Majorana mass  $|\Delta|.$

## Zero Mode

In the presence of a single vortex order parameter  $\Delta(\mathbf{r}) = v(r)e^{i\theta}$  there exists a zero-energy (static) isolated mode

(Fu & Kane, *PRL* **100**, 096407 (08); Rossi & RJ *NPB* **190**, 681 (81))

Majorana field expansion:

$$\Psi = \dots\dots\dots + a \Psi_0$$

$E \neq 0$  modes

where zero mode operator  $a$  satisfies

$$\{a, a^\dagger\} = 1, a^\dagger = a \Rightarrow a^2 = 1/2$$